



# Evaluate the effects of trauma in adults with the gold standard

The TSI-2 evaluates posttraumatic stress and other psychological sequelae from traumatic events including sexual and physical assault, partner violence, combat, torture, motor vehicle accidents, medical trauma, mass casualty events, traumatic loss, and childhood abuse or neglect.



Learn more at [parinc.com/TSI2](https://parinc.com/TSI2)

**Purpose** Evaluates acute and chronic posttraumatic symptomatology

**Age** 18 years and older

**Format** Paper and pencil; online administration and scoring via PARiConnect

**Time** 20 minutes to administer;  
20 minutes to score

**Qual** B

**PARiConnect**

**PAR** | TRAINING

## A broadband measure of posttraumatic symptomatology

- Three scales (Insecure Attachment, Somatic Preoccupations, and Suicidality), several subscales, and four summary factors (Self-Disturbance, Posttraumatic Stress, Externalization, and Somatization) are new or have been significantly reconfigured.
- Items assess clients' tendencies to deny symptoms that are commonly endorsed, to overendorse unusual or bizarre symptoms, or to respond in an inconsistent or random manner. Also addresses malingering.
- Available in Spanish.
- A 136-item carbonless form assesses a wide range of potentially complex symptomatology ranging from PTSD, dissociation, and somatization to insecure attachment styles, impaired self-capacities, and dysfunctional behaviors.
- Eight critical items help identify issues that potentially represent severe psychological disturbance or danger to the respondent or others.
- Features an alternate form (TSI-2-A) that does not contain sexual symptom items.
- The validation sample consisted of five nonoverlapping groups: Combat veterans, individuals with borderline personality disorder, sexual abuse victims, victims of domestic violence, and incarcerated women.
- Convenient administration and scoring on PARiConnect, our online assessment platform.