

# References

- Alexander, G. E., DeLong, M. R., & Strick, P. L. (1986). Parallel organization of functionally segregated circuits linking basal ganglia and cortex. *Annual Review of Neuroscience*, 9, 357-381.
- American Educational Research Association, American Psychological Association, & the National Council on Measurement in Education. (1999). *Standards for educational and psychological testing*. Washington, DC: American Educational Research Association.
- American Psychiatric Association. (1994). *Diagnostic and statistical manual of mental disorders* (4th ed.). Washington, DC: Author.
- Beck, A. T., & Steer, R. A. (1990). *Beck Anxiety Inventory (BAI) manual*. San Antonio, TX: The Psychological Corporation.
- Beck, A. T., Steer, R. A., & Brown, G. K. (1996). *Beck Depression Inventory-II (BDI-II) manual*. San Antonio, TX: The Psychological Corporation.
- Benedict, R. H. B. (1997). *Brief Visuospatial Memory Test-Revised (BVMT-R): Professional manual*. Odessa, FL: Psychological Assessment Resources.
- Benton, A. L., & Hamsher, K. deS. (1976). *Multilingual Aphasia Examination: Manual*. Iowa City: University of Iowa. (Rev. ed., 1978).
- Benton, A. L., Hannay, H. J., & Varney, N. R. (1975). Visual perception of line direction in patients with unilateral brain disease. *Neurology*, 25, 907-910.
- Bigler, E. D. (2001). Frontal lobe pathology and anti-social personality disorder. *Archives of General Psychiatry*, 58(6), 609-611.
- Borod, J. C., Goodglass, H., & Kaplan, E. (1980). Normative data on the Boston Diagnostic Aphasia Examination, Parietal Lobe Battery, and the Boston Naming Test. *Journal of Clinical Neuropsychology*, 2, 209-216.
- Boyle, P. A., Grace, J., Zawacki, T. M., Ott, B. R., & Stout, J. C. (2001). *Frontal behavior in neurodegenerative and acute neurological disorders*. Manuscript submitted for publication.
- Brandt, J. (1991). The Hopkins Verbal Learning Test: Development of a new verbal memory test with six equivalent forms. *The Clinical Neuropsychologist*, 5, 125-142.
- Breaden, C. E., Hoffman, K. M., & Cannon, T. D. (2001). The neuropsychology and neuroanatomy of bipolar affective disorder: A critical review. *Bipolar Disorder*, 3, 106-150.
- Brown, G., Chadwick, O., Shaffer, D., Rutter, M., & Traub, M. (1981). A prospective study of children with head injuries: III. Psychiatric sequelae. *Psychological Medicine*, 11(1), 63-78.
- Butcher, J. N., Dahlstrom, W. G., Graham, J. R., Tellegen, A., & Kaemmer, B. (1989). *Minnesota Multiphasic Personality Inventory-2 (MMPI-2): Manual for administration and scoring*. Minneapolis: University of Minnesota Press.
- Cahn-Weiner, D., Grace, J., Ott, B. R., Fernandez, H., & Friedman, J. F. (in press). Cognitive and behavioral features of Alzheimer's and Parkinson's disease. *Neuropsychiatry, Neuropsychology, and Behavioral Neurology*.
- Cahn-Weiner, D. A., Malloy, P. F., Boyle, P. A., Marran, M., & Salloway, S. (2000). Prediction of functional status from neuropsychological tests in community-dwelling elderly individuals. *Clinical Neuropsychologist*, 14(2), 187-195.
- Cicerone, K., & Tanenbaum, L. N. (1997). Disturbance of social cognition after traumatic orbitofrontal brain injury. *Archives of Clinical Neuropsychology*, 12(2), 173-188.
- Cronbach, L. J. (1951). Coefficient alpha and the internal structure of tests. *Psychometrika*, 16, 297-334.

- Cummings, J. L. (1993). Frontal-subcortical circuits and human behavior. *Archives of Neurology*, 50(8), 873-880.
- Cummings, J. L., Mega, M., Gray, K., Rosenberg-Thompson, S., Carusi, D. A., & Gornbein, J. (1994). The Neuropsychiatric Inventory: Comprehensive assessment of psychopathology in dementia. *Neurology*, 44(12), 2308-2314.
- Damasio, A. R., & Van Hoesen, G. (1983). Emotional disturbances associated with focal lesions of the limbic frontal lobe. In K. M. Heilman & P. Satz (Eds.), *Neuropsychology of human emotion* (pp. 85-110). New York: Guilford.
- Deckel, A. W., Hesselbrock, V., & Bauer, L. (1996). Antisocial personality disorder, childhood delinquency, and frontal brain functioning: EEG and neuropsychological findings. *Journal of Clinical Psychology*, 52(6), 639-650.
- Dinn, W. M., & Harris, C. L. (2000). Neurocognitive function in antisocial personality disorder. *Psychiatry Research*, 97(2-3), 173-190.
- Eslinger, P. J., & Damasio, A. R. (1985). Severe disturbance of higher cognition after bilateral frontal lobe ablation: Patient EVR. *Neurology*, 35(12), 1731-1741.
- Flaherty, J. H., Miller, D. K., & Coe, R. M. (1992). Impact on caregivers of supporting urinary function in noninstitutionalized, chronically ill seniors. *Gerontologist*, 32(4), 541-545.
- Folstein, M. F., Folstein, S. E., & McHugh, P. R. (2001). *Mini-Mental State Examination*. Odessa, FL: Psychological Assessment Resources.
- Gioia, G. A., Isquith, P. K., Guy, S. C., & Kenworthy, L. (2000). *BRIEF: Behavior Rating Inventory of Executive Function professional manual*. Odessa, FL: Psychological Assessment Resources.
- Goldman-Rakic, P. S. (1995). Architecture of the prefrontal cortex and the central executive. In J. Grafman, K. F. Holyoak, & F. Boller (Eds.). Structure and functions of the human prefrontal cortex. *Annals of the New York Academy of Science*, 179, 71-83.
- Gorenstein, E. E. (1982). Frontal lobe functions in psychopaths. *Journal of Abnormal Psychology*, 91(5), 368-379.
- Gorsuch, R. L. (1997). Exploratory factor analysis: Its role in item analysis. *Journal of Personality Assessment*, 68(3), 532-560.
- Grace, J., Boyle, P., Zawacki, T., & Stout, J. C. (2000). Frontal behavior change in chronic and acute neurological disorders. *Journal of Neuropsychiatry and Clinical Neurosciences*, 12(1), 146.
- Grace, J., Malloy, P. F., & Stout, J. C. (1997). Quantitative assessment of frontal apathy: A case study utilizing the Frontal Lobe Personality Scale. *Journal of Neuropsychiatry and Clinical Neurosciences*, 9, 672.
- Grace, J., Malloy, P. F., Stout, J. C., & the National Association of Neuropsychology Research Consortium. (1997). Assessing frontal behavioral syndromes: Reliability and validity of the Frontal Lobe Personality Scale. *Archives of Clinical Neuropsychology*, 12, 327.
- Grace, J., Stout, J. C., & Dupont, R. (1997). [Comparison of frontal lesion and psychiatric patient groups on the Frontal Systems Behavior Scale.] Unpublished raw data.
- Grace, J., Stout, J. C., & Malloy, P. F. (1999). Assessing frontal lobe behavioral syndromes with the Frontal Lobe Personality Scale. *Assessment*, 6(3), 269-284.
- Grant, D. A., & Berg, E. A. (1948). A behavioral analysis of degree of reinforcement and ease of shifting to new responses in a Weigl-type card sorting problem. *Journal of Experimental Psychology*, 34, 404-411.
- Grober, E., & Sliwinski, M. (1991). Development and validation of a model for estimating premorbid verbal intelligence in the elderly. *Journal of Clinical & Experimental Neuropsychology*, 13(6), 933-949.
- Grodzinsky, G. M., & Barkley, R. A. (1999). Predictive power of frontal lobe tests in the diagnosis of attention deficit hyperactivity disorder. *Clinical Neuropsychology*, 13(1), 12-21.
- Gustafsson, P., Thernlund, G., Ryding, E., Rosen, I., & Cederblad, M. (2000). Associations between cerebral blood-flow measured by single photon emission computed tomography (SPECT), electroencephalogram (EEG), behaviour symptoms, cognition and neurological soft signs in children with attention-deficit hyperactivity disorder (ADHD). *Acta Paediatrica*, 89(7), 830-835.

- Hachinski, V. C., Illiff, L. D., Zilhka, E., Du Boulay, G. H., McAllister, V. L., Marshall, J., et al. (1975). Cerebral blood flow in dementia. *Archives of Neurology*, 32(9), 632-637.
- Harlow, J. (1848). Passage of an iron bar through the head. *Boston Medical and Surgical Journal*, 39, 389-393.
- Hendren, R. L., De Backer, I., & Pandina, G. J. (2000). Review of neuroimaging studies of child and adolescent psychiatric disorders from the past 10 years. *Journal of the American Academy of Child Adolescent Psychiatry*, 39(7), 815-828.
- Hoehn, M. M., & Yahr, M. D. (1967). Parkinsonism: Onset, progression, and mortality. *Neurology*, 17, 427-442.
- Hooper, H. E. (1983). *Hooper Visual Organization Test (VOT)*. Los Angeles: Western Psychological Services.
- Hughes, C. P., Berg, L., Danzinger, W. L., Coben, L. A., & Martin, R. L. (1982). A new clinical scale for the staging of dementia. *British Journal of Psychiatry*, 140, 566-572.
- Jones-Gotman, M., & Zatorre, R. J. (1988). Olfactory identification deficits in patients with focal cerebral excision. *Neuropsychologia*, 26(3), 387-400.
- Jurica, P. J., Leitten, C. L., & Mattis, S. (2001). *DRS-2: Dementia Rating Scale-2 professional manual*. Odessa, FL: Psychological Assessment Resources.
- Kaplan, E. F., Goodglass, H., & Weintraub, S. (1983). *The Boston Naming Test* (2nd ed.). Philadelphia: Lea & Febiger.
- Kertesz, A., Davidson, W., & Fox, H. (1997). Frontal Behavioral Inventory: Diagnostic criteria for frontal lobe dementia. *Canadian Journal of Neurological Science*, 24(1), 29-36.
- Kertesz, A., Nadkarni, N., Davidson, W., & Thomas, A. W. (2000). The Frontal Behavioral Inventory in the differential diagnosis of frontotemporal dementia. *Journal of the International Neuropsychological Society*, 6(4), 460-468.
- Kiernan, R. J., Mueller, J., Langston, J. W., & VanDyke, C. (1987). The Neurobehavioral Cognitive Status Examination. *Annals of Internal Medicine*, 107, 481-485.
- Kuruoglu, A. C., Arikan, Z., Vural, G., Karatas, M., Arac, M., & Isik, E. (1996). Single photon emission computerised tomography in chronic alcoholism. Antisocial personality disorder may be associated with decreased frontal perfusion. *British Journal of Psychiatry*, 169(3), 348-354.
- Larson, E. R., Shear, P. K., Krikorian, R., & Strakowski, S. M. (2001). Frontal lobe mediated behavior and mood state in bipolar disorder. *Clinical Neuropsychologist*, 15, 273.
- Levin, H. S., Eisenberg, H. M., & Benton, A. L. (1991). *Frontal lobe function and dysfunction*. New York: Oxford University Press.
- Levy, M. L., Cummings, J. L., Fairbanks, L. A., Masterman, D., Miller, B. L., Craig, A. H., et al. (1998). Apathy is not depression. *Journal of Neuropsychiatry*, 10, 314-319.
- Lhermitte, F., Pillon, B., & Serdaru, M. (1986). Human autonomy and the frontal lobes. Part I: Imitation and utilization behavior: A neuropsychological study of 75 patients. *Annals of Neurology*, 19(4), 326-334.
- Lichter, D. G., & Cummings, J. L. (Eds.). (2001). *Frontal-subcortical circuits in psychiatric and neuropsychiatric disorders*. New York: Guilford.
- Lueger, R. J., & Gill, K. J. (1990). Frontal-lobe cognitive dysfunction in conduct disorder adolescents. *Journal of Clinical Psychology*, 46(6), 696-706.
- The Lund and Manchester Groups. (1994). Clinical and neuropathological criteria for frontotemporal dementia. *Journal of Neurology, Neurosurgery, and Psychiatry*, 57(4), 416-418.
- Malloy, P. F., & Aloia, M. (1998). Frontal lobe dysfunction in traumatic brain injury. *Seminars in Clinical Neuropsychiatry*, 3(3), 186-194.
- Malloy, P., Bahrle, A., Duffy, J., & Cimino, C. (1993). The orbitomedial frontal syndrome. *Archives of Clinical Neuropsychology*, 8(3), 185-201.
- Malloy, P. F., & Duffy, J. (1994). The frontal lobes in neuropsychiatric disorders. In F. Boller & J. Grafman (Eds.). *Handbook of neuropsychology*: Vol. 9 (pp. 203-232). New York: Elsevier Science B.V.

- Malloy, P., Noel, N., Longabaugh, R., & Beattie, M. (1990). Determinants of neuropsychological impairment in antisocial substance abusers. *Addictive Behaviors*, 15(5), 431-438.
- Mattis, S. (1988). *Dementia Rating Scale: Professional manual*. Odessa, FL: Psychological Assessment Resources.
- McKhann, G., Drachman, D., Folstein, M., Katzman, R., Price, D., & Stadlan, E. M. (1984). Clinical diagnosis of Alzheimer's disease: Report of the NINCDS-ADRDA work group under the auspices of the Department of Health and Human Services Task Force on Alzheimer's disease. *Neurology*, 34(7), 939-944.
- Mega, M. S., & Cummings, J. L. (1994). Frontal-subcortical circuits and neuropsychiatric disorders. *Journal of Neuropsychiatry and Clinical Neurosciences*, 6(4), 358-370.
- Mesulam, M. (2000). *Principles of behavioral and cognitive neurology*. New York: Oxford University Press.
- Mesulam, M.-M. (1985). *Principles of behavioral neurology*. Philadelphia: F. A. Davis.
- Mindus, P., Edman, G., & Andreewitch, S. (1999). A prospective long-term study of personality traits in patients with intractable obsessional illness treated by capsulotomy. *Acta Psychiatrica Scandinavica*, 99(1), 40-50.
- Morris, J. C. (1993). The Clinical Dementia Rating (CDR): Current version and scoring rules. *Neurology*, 43, 2412-2414.
- Morris, J. C., Heyman, A., Mohs, R. C., Hughes, J. P., Van Belle, G., Fillenbaum, G., Mellits, E. D., Clark, C., & the CERAD Investigators. (1989). The Consortium to Establish a Registry for Alzheimer's Disease (CERAD)/Part I. Clinical and neuropsychological assessment of Alzheimer's disease. *Neurology*, 39, 1159-1165.
- Moser, D. J., Cohen, R. A., Malloy, P. F., Stone, W. M., & Rogg, J. M. (1998). Reduplicative paramnesia: Longitudinal neurobehavioral and neuroimaging analysis. *Journal of Geriatric Psychiatry & Neurology*, 11(4), 174-180.
- Nagaratnam, N., Lewis-Jones, M., Scott, D., & Palazzi, L. (1998). Behavioral and psychiatric manifestations in dementia patients in a community: Caregiver burden and outcome. *Alzheimer's Disease and Associated Disorders*, 12, 330-334.
- Neary, D., Snowden, J. S., Gustafson, L., Passant, U., Stuss, D., Black, S., et al. (1998). Frontotemporal lobar degeneration: A consensus on clinical diagnostic criteria. *Neurology*, 51(6), 1546-1554.
- Norton, L. E., Malloy, P. F., & Salloway, S. (2001). The impact of behavioral symptoms on activities of daily living in patients with dementia. *American Journal of Geriatric Psychiatry*, 9(1), 41-48.
- Paulsen, J. S., Ready, R. E., Stout, J. C., Salmon, D. P., Thal, L. J., Grant, I., et al. (2000). Neurobehaviors and psychotic symptoms in Alzheimer's disease. *Journal of the International Neuropsychological Society*, 6(7), 815-820.
- Paulsen, J. S., Stout, J. C., DelaPena, J., Romero, R., Tawfik-Reedy, Z., Swenson, M. R., et al. (1996). Frontal behavioral syndromes in cortical and subcortical dementia. *Assessment*, 3, 327-337.
- The Psychological Corporation. (1997). *WAIS-III/WMS-III technical manual*. San Antonio, TX: Author.
- The Psychological Corporation. (1999). *Wechsler Abbreviated Scale of Intelligence (WASI) manual*. San Antonio, TX: Author.
- Ready, R., Grace, J., Ott, B. R., & Cahn-Weiner, D. (2001). Behavior change in mild cognitive impairment and dementia. *Archives of Clinical Neuropsychology*, 16(8), 712.
- Reise, S., Waller, N., & Comrey, A. (2000). Factor analysis and scale revision. *Psychological Assessment*, 12(3), 287-297.
- Reitan, R. (1958). Validity of the Trail Making Test as an indication of organic brain damage. *Perceptual and Motor Skills*, 8, 271-276.
- Rubia, K., Overmeyer, S., Taylor, E., Brammer, M., Williams, S. C., Simmons, A., et al. (1999). Hypofrontality in attention deficit hyperactivity disorder during higher-order motor control: A study with functional MRI. *American Journal of Psychiatry*, 156(6), 891-896.

- Rymer, S., Salloway, S., Norton, L., & Malloy, P. F. (2001). *Frontal behavioral disturbance and caregiver burden*. Manuscript submitted for publication.
- Saint-Cyr, J. A., & Trepanier, L. L. (2000). Neuropsychologic assessment of patients for movement disorder surgery. *Movement Disorders*, 15(5), 771-783.
- Saint-Cyr, J. A., Trepanier, L. L., Kumar, R., Lozano, A. M., & Lang, A. E. (2000). Neuropsychological consequences of chronic bilateral stimulation of the subthalamic nucleus in Parkinson's disease. *Brain*, 123, 2091-2108.
- Sano, M., Devanand, D. P., Richards, M., Miller, L. W., Marder, K., Bell, K., et al. (1995). A standardized technique for establishing onset and duration of symptoms of Alzheimer's disease. *Archives of Neurology*, 52(10), 961-966.
- Selemon, L. D., & Goldman-Rakic, P. S. (1985). Longitudinal topography and interdigitation of corticostriatal projections in the rhesus monkey. *Journal of Neuroscience*, 5(3), 776-794.
- Sheikh, J. I., & Yesavage, J. A. (1986). Geriatric Depression Scale (GDS): Recent evidence and development of a shorter version. In T. L. Brink (Ed.), *Clinical gerontology: A guide to assessment and intervention* (pp. 165-173). New York: Haworth Press.
- Solomon, D. A., & Malloy, P. F. (1992). Alcohol, head injury, and neuropsychological function. *Neuropsychology Review*, 3(3), 249-280.
- Solyom, L., Turnbull, I. M., & Wilensky, M. (1987). A case of self-inflicted leucotomy. *British Journal of Psychiatry*, 151, 855-857.
- Spreen, O., & Strauss, E. (1998). Language tests. In *A Compendium of Neuropsychological Tests: Administration, Norms and Commentary* (2nd ed., pp. 447-459). New York: Oxford University Press.
- Stout, J. C., Ready, R., Grace, J., Malloy, P. F., & Paulsen, J. (2001). *Factor analysis of the Frontal Systems Behavior Scale*. Manuscript submitted for publication.
- Stout, J. C., Wyman, M. F., Peavy, G. M., & Salmon, D. P. (2001). *Frontal neuropsychiatric symptoms in probable Alzheimer's disease*. Manuscript submitted for publication.
- Stroop, J. R. (1935). Studies of interference in serial verbal reactions. *Journal of Experimental Psychology*, 18, 643-662.
- Stuss, D., & Benson, D. (1986). *The frontal lobes*. New York: Raven Press.
- Stuss, D. T., & Benson, D. F. (1984). Neuropsychological studies of the frontal lobes. *Psychological Bulletin*, 95(1), 3-28.
- Trepanier, L. L., Kumar, R., Lozano, A. M., Lang, A. E., & Saint-Cyr, J. A. (2000). Neuropsychological outcome of GPi pallidotomy and GPi or STN deep brain stimulation in Parkinson's disease. *Brain and Cognition*, 42, 324-347.
- Trepanier, L. L., Saint-Cyr, J. A., Lozano, A. M., & Lang, A. E. (1998). Neuropsychological consequences of posteroverentral pallidotomy for the treatment of Parkinson's disease. *Neurology*, 51(1), 207-215.
- Wechsler, D. (1945). A standardized memory scale for clinical use. *Journal of Psychology*, 19, 87-95.
- Wechsler, D. (1955). *Wechsler Adult Intelligence Scale (WAIS) manual*. New York: The Psychological Corporation.
- Wechsler, D. (1981). *Wechsler Adult Intelligence Scale-Revised (WAIS-R) manual*. New York: The Psychological Corporation.
- Welsh, K. A., Butters, N., Mohs, R. C., Beekly, D., Edland, S., Fillenbaum, G., et al. (1994). The Consortium to Establish a Registry for Alzheimer's Disease (CERAD). Part V. A normative study of the neuropsychological battery. *Neurology*, 44(4), 609-614.
- Zarit, S., Reever, K., & Bach-Peterson, B. (1980). Relatives of the impaired elderly: Correlates of feelings of burden. *The Gerontologist*, 20, 649-655.
- Zawacki, T. M., Grace, J., Paul, R. H., Moser, D. J., Ott, B. R., Gordon, N., et al. (in press). Impact of apathy on the prediction of functional abilities of vascular dementia patients. *Journal of Neuropsychiatry and Clinical Neurosciences*.